

On k -Transversals

RON AHARONI

Department of Mathematics, Technion–I.I.T., Haifa 32000, Israel

AND

PÉTER KOMJÁTH

*Department of Algebra and Number Theory,
R. Eötvös University, P.O.B. 323, 1445 Budapest, Hungary*

Communicated by the Managing Editors

Received July 29, 1983; revised June 18, 1986

A k -transversal of a family of sets \mathcal{H} is a function $g: \mathcal{H} \rightarrow [\bigcup \mathcal{H}]^{k+1}$ satisfying $|\bigcup g[\mathcal{F}]| \geq |\mathcal{F}| + k$ for every nonempty subfamily \mathcal{F} of \mathcal{H} and $g(H) \subseteq H$ for every $H \in \mathcal{H}$. It is proved that a (possibly infinite) family \mathcal{H} has a k -transversal if and only if the family $(H \setminus S: H \in \mathcal{H})$ has a transversal for every $S \in [\bigcup \mathcal{H}]^k$. One corollary is that a family of infinite sets has a k -transversal if and only if it has a transversal. It is also shown that if $l \leq k$ and $f(H) \in [H]^l$ for each $H \in \mathcal{H}$ and \mathcal{H} has a k -transversal, then it has a k -transversal g such that $g(H) \supseteq f(H)$ for every $H \in \mathcal{H}$. © 1987 Academic Press, Inc.

I. INTRODUCTION AND STATEMENT OF THE RESULTS

Throughout this paper \mathcal{H} denotes a (possibly infinite) nonempty family of sets and V denotes $\bigcup \mathcal{H}$. Also, k denotes any natural number. A function $g: \mathcal{H} \rightarrow [V]^{k+1}$ (the set of subsets of V of size $k+1$) is called a k -transversal if $|\bigcup g[\mathcal{F}]| \geq |\mathcal{F}| + k$ for every nonempty subfamily \mathcal{F} of \mathcal{H} and $g(H) \subseteq H$ for every $H \in \mathcal{H}$. This is a generalisation of the notion of a “transversal,” which is just a 0-transversal. (However, “transversals” have below distinct notation in that they are regarded as functions into V rather than into $[V]^1$.) It was introduced by Lovász [5], who proved the following extension of Hall’s theorem [3]: a finite family \mathcal{H} has a k -transversal if and only if $|\bigcup \mathcal{F}| \geq |\mathcal{F}| + k$ for every nonempty subfamily \mathcal{F} of \mathcal{H} . The main result of this paper is one which clearly implies Lovász’ characterisation, and which is valid also for infinite families. For any subset S of V denote by $\mathcal{H} - S$ the family $(H \setminus S: H \in \mathcal{H})$.

THEOREM 1. *\mathcal{H} has a k -transversal if and only if $|V| \geq k$ and $\mathcal{H} - S$ has a transversal for every $S \in [V]^k$.*

A string f in V is an injection from some ordinal $\alpha = \text{dom } f$ into V . For $A \subseteq V$ define $D(A) = (H \in \mathcal{H} : H \subseteq A)$. For any set T write $\|T\| = |T|$ if T is finite and $\|T\| = \infty$ if T is infinite. We adopt the conventions $\infty - \infty = \infty$, and $r + \infty = \infty$, $r - \infty = -\infty$ for any integer r . Nash-Williams [6] defined the following function q on strings in V :

$$q(\emptyset) = -\|D(\emptyset)\|;$$

if $\alpha = \text{dom } f = \beta + 1$ then

$$q(f) = q(f \restriction \beta) + 1 - \|D(f[\alpha]) \setminus D(f[\beta])\|;$$

if α is a limit ordinal then

$$q(f) = \liminf_{\beta < \alpha} q(f \restriction \beta) - \left\| D(f[\alpha]) \setminus \bigcup_{\beta < \alpha} D(f[\beta]) \right\|.$$

It is easy to show that if \mathcal{H} has a transversal then $q(f) \geq 0$ holds for every string f . Nash-Williams proved that for countable \mathcal{H} this condition implies the existence of a transversal. In [1] this was shown to hold also if $\mathcal{H} \setminus \mathcal{H}'$ has a transversal for some countable subfamily \mathcal{H}' . As a corollary of Theorem 1 there follows:

COROLLARY 1a. *The family \mathcal{H} with $|V| \geq k$ has a k -transversal if and only if it has a transversal and $q(f) \geq k$ for every string f in V with $|\text{dom } f| \geq k$.*

A subfamily \mathcal{C} of \mathcal{H} is *critical* if it has a transversal and $g[\mathcal{C}] = \bigcup \mathcal{C}$ for every transversal g of \mathcal{C} . Podewski and Steffens [7] proved that if \mathcal{H} is countable, then it has a transversal if and only if it does not contain a critical subfamily \mathcal{C} , such that $H \subseteq \bigcup \mathcal{C}$ for some $H \in \mathcal{H} \setminus \mathcal{C}$.

Another consequence of Theorem 1 is

COROLLARY 1b. *\mathcal{H} has a 1-transversal if and only if it has a transversal and it contains no nonempty critical subfamily.*

For general k the corresponding result is

COROLLARY 1c. *\mathcal{H} has a k -transversal if and only if it has a transversal and it does not contain a subfamily \mathcal{C} such that $|\bigcup \mathcal{C} \setminus f[\mathcal{C}]| < k$ for every transversal f of \mathcal{C} .*

One of the main tools for the proof of Theorem 1 is a result which is of interest in itself:

THEOREM 2. *If $l \leq k$ and $f: \mathcal{H} \rightarrow [V]^l$ satisfies $f(H) \subseteq H$ and \mathcal{H} has a k -transversal, then \mathcal{H} has a k -transversal g such that $g(H) \supseteq f(H)$ for every $H \in \mathcal{H}$.*

A family \mathcal{H} is *two-colorable* if there exists a coloring of V in two colors such that every $H \in \mathcal{H}$ contains vertices of both colors. If \mathcal{H} has a 1-transversal, then it is two-colorable, since a 1-transversal can be considered as a forest (its 2-sets being taken as edges of a graph on V); and a forest is clearly 2-colorable. In [4] it was proved that if all members of \mathcal{H} are infinite and \mathcal{H} has a transversal, then it is two-colorable. Here we strengthen this to

THEOREM 3. *If all members of \mathcal{H} are infinite then \mathcal{H} has a k -transversal if and only if it has a transversal.*

Theorem 3 was suggested to us by E. Milner, and it was the attempt to prove it which started the investigations of this paper.

II. PROOFS

LEMMA 1. *Suppose that $k > 0$ and that g is a k -transversal of \mathcal{H} . If $F \in \mathcal{H}$ and $x \in F \setminus g(F)$ and a and b are distinct members of $g(F)$ then for $y = a$ or for $y = b$ there holds: the function g' defined by $g'(F) = g(F) \setminus \{y\} \cup \{x\}$ and $g'(I) = g(I)$ for $I \neq F$ is a k -transversal.*

Proof. Suppose that neither a nor b can serve in the role of y in the lemma. Then there exist finite subfamilies \mathcal{I} and \mathcal{J} of $\mathcal{H} \setminus \{F\}$ such that

$$|\bigcup g[\mathcal{I}] \cup (g(F) \setminus \{a\} \cup \{x\})| < |\mathcal{I}| + 1 + k \quad (1)$$

and

$$|\bigcup g[\mathcal{J}] \cup (g(F) \setminus \{b\} \cup \{x\})| < |\mathcal{J}| + 1 + k. \quad (2)$$

Let $\mathcal{K} = \mathcal{I} \cup \mathcal{J} \cup \{F\}$. Suppose first that $\mathcal{I} \cap \mathcal{J} \neq \emptyset$. Then, since g is a k -transversal, $|\bigcup g[\mathcal{I} \cap \mathcal{J}]| \geq |\mathcal{I} \cap \mathcal{J}| + k$. By (1) and (2) this entails

$$|\bigcup g[\mathcal{K}]| \leq |\mathcal{I}| + k + |\mathcal{J}| + k - (|\mathcal{I} \cap \mathcal{J}| + k) = |\mathcal{K}| - 1 + k,$$

contradicting the fact that g is a k -transversal. Thus we may assume that $\mathcal{I} \cap \mathcal{J} = \emptyset$.

Since $|g(F)| = k + 1$ we have

$$\begin{aligned} & |(\bigcup g[\mathcal{I}] \cup (g(F) \setminus \{a\} \cup \{x\})) \cap (\bigcup g[\mathcal{J}] \cup (g(F) \setminus \{b\} \cup \{x\}))| \\ & \geq |g(F) \setminus \{a, b\} \cup \{x\}| = k. \end{aligned}$$

Therefore, by (1) and (2),

$$|\bigcup g[\mathcal{H}]| \leq |\mathcal{I}| + k + |\mathcal{J}| + k - k = |\mathcal{H}| - 1 + k,$$

again contradicting the fact that g is a k -transversal.

COROLLARY. *If g is a k -transversal of \mathcal{H} and $H \in \mathcal{H}$ and F is a subset of H and $|F| \leq k$, then there exists a k -transversal \bar{g} of \mathcal{H} such that $\bar{g}(H) \supseteq F$ and $\bar{g}(I) = g(I)$ for each $I \neq H$.*

Proof. It suffices to show that if $F \not\subseteq g(H)$ then it is possible to replace $g(H)$ by $g(H) \setminus \{y\} \cup \{x\}$ for some $y \in g(H) \setminus F$ and $x \in F \setminus g(H)$ while keeping g a k -transversal. But if $F \not\subseteq g(H)$ then there exists $x \in F \setminus g(H)$ and since $|F| \leq |g(H)| - 1$ there exist distinct $a, b \in g(H) \setminus F$, and one can apply the lemma.

The proof of Theorem 2 now follows easily. Let h be a k -transversal of \mathcal{H} . We well order \mathcal{H} as $(H_\theta : \theta < |\mathcal{H}|)$, and one by one replace $h(H_\theta)$ by a $k+1$ -subset of H_θ which contains $f(H_\theta)$, while keeping h a k -transversal. The resulting function g is clearly a k -transversal, since it suffices to check the condition $|\bigcup g[\mathcal{F}]| \geq |\mathcal{F}| + k$ for finite subfamilies of \mathcal{H} (and this is also the reason why in limit steps of the definition of g we are still left with a k -transversal).

We shall need Theorem 2 only in the case that f is a transversal.

LEMMA 2. *If \mathcal{H} has a transversal f and it can be decomposed as $\mathcal{H} = \bigcup_{\alpha < \zeta} \mathcal{H}_\alpha$ so that each \mathcal{H}_α has a k -transversal and $f[\mathcal{H}_\alpha] \cap \bigcup \mathcal{H}_\beta = \emptyset$ whenever $\alpha > \beta$, then it has a k -transversal.*

Proof. By Theorem 2 each \mathcal{H}_α has a k -transversal g_α such that $f(H) \in g_\alpha(H)$ for $H \in \mathcal{H}_\alpha$. Define: $g = \bigcup_{\alpha < \zeta} g_\alpha$. Let \mathcal{G} be a finite subfamily of \mathcal{H} . Let α be the first ordinal such that $\mathcal{G} \cap \mathcal{H}_\alpha \neq \emptyset$. Then

$$\begin{aligned} |\bigcup g[\mathcal{G}]| &\geq |\bigcup g_\alpha[\mathcal{G} \cap \mathcal{H}_\alpha]| + |f[\mathcal{G} \setminus \mathcal{H}_\alpha]| \\ &\geq |\mathcal{G} \cap \mathcal{H}_\alpha| + k + |\mathcal{G} \setminus \mathcal{H}_\alpha| = |\mathcal{G}| + k. \end{aligned}$$

Proof of Theorem 1. Assume first that \mathcal{H} has a k -transversal g . Let $S \subseteq V$ satisfy $|S| \leq k$. For each $H \in \mathcal{H}$ define $g'(H) = g(H) \setminus S$. Then $|\bigcup g'[\mathcal{F}]| \geq |\mathcal{F}|$ for any $\mathcal{F} \subseteq \mathcal{H}$. Since the sets $g'(H)$ are finite, it follows by Hall's theorem [2] that the family $(g'(H) : H \in \mathcal{H})$ has a transversal. Since $g'(H) \subseteq H \setminus S$, so does $\mathcal{H} - S$.

For the proof of the reverse direction of Theorem 1, we need first the following strengthening of the finite case of the theorem:

LEMMA 3. Let $\mathcal{H} = (H_0, \dots, H_n)$ be a finite family. For $0 \leq i \leq n$ write $\mathcal{H}_i = (H_j: j \geq i)$. If for each $0 \leq i \leq n$ there exists a set $A_i \in [H_i]^k$ such that $\mathcal{H}_i - A_i$ has a transversal f_i , then \mathcal{H} has a k -transversal.

Proof. By induction on n . For $n=0$ the lemma is clear. Let now $n > 0$. By the induction hypothesis there exists a k -transversal g of \mathcal{H}_1 . By Theorem 2 we may assume that $f_0(H_i) \in g(H_i)$ for $1 \leq i \leq n$. Extend g to a function on \mathcal{H} by defining $g(H_0) = A_0 \cup \{f_0(H_0)\}$. Applying Lemma 2 with $f, \mathcal{H}, \zeta, \mathcal{H}_1, \mathcal{H}_0$ replaced respectively by $f^*, (g(H_i): 0 \leq i \leq n), 2, (g(H_i): 1 \leq i \leq n)$ and the family whose sole member is $g(H_0)$, where $f^*(g(H_i)) = f_0(H_i - A_0)$, we see that g is a k -transversal.

Lemma 3 clearly yields Theorem 1 for finite \mathcal{H} . Consider next the countable case, i.e., when $\mathcal{H} = (H_i: i < \omega)$. For each $i < \omega$ choose $A_i \in [H_i]^k$. By the assumption of the theorem there exists a transversal f_i of $\mathcal{H} - A_i$. For each $i < \omega$ let $H'_i = A_i \cup \{f_j(H_i \setminus A_j): j \leq i\}$. For $m < \omega$ the family $(H'_0, H'_1, \dots, H'_m)$ satisfies the conditions of Lemma 3, and hence has a k -transversal. But each H'_i is finite, and by compactness it follows that the family $(H'_i: i < \omega)$ has a k -transversal. Since $H'_i \subseteq H_i$, so does \mathcal{H} .

Consider now the case in which \mathcal{H} is of general cardinality. By the assumption of the theorem, there exists a transversal f of \mathcal{H} . We shall show that it is possible to split \mathcal{H} as

$$\mathcal{H} = \bigcup_{\alpha < \zeta} \mathcal{H}_\alpha \quad \text{and } V \text{ as } V = \bigcup_{\alpha < \zeta} V_\alpha \quad \text{so that:}$$

- (i) $\mathcal{H}_\alpha \cap \mathcal{H}_\beta = V_\alpha \cap V_\beta = \emptyset$ whenever $\alpha \neq \beta$;
- (ii) $f[\mathcal{H}_\alpha] \subseteq V_\alpha$; and
- (iii) each \mathcal{H}_α has a k -transversal contained in $\bigcup_{\beta \leq \alpha} V_\beta$.

By Lemma 2 it will follow then that \mathcal{H} has a k -transversal.

We well order \mathcal{H} as $(H_\theta: \theta < \mu)$. Choose $A_0 \in [H_0]^k$ and let f_1 be a transversal of \mathcal{H} with $f_1[\mathcal{H}] \cap A_0 = \emptyset$. For any transversal h of \mathcal{H} write $E(h) = \{\{H, h(H)\}: H \in \mathcal{H}\}$. Let Γ_1 be a bipartite graph whose sides are \mathcal{H} and V , and whose edge set is $E_1 = E(f) \cup E(f_1)$. Let C_1 be the connected component of Γ_1 containing H_0 . Clearly C_1 is countable. Choose a set A_1 in $[V]^k$ different from A_0 . Let f_2 be a transversal of \mathcal{H} with $f_2[\mathcal{H}] \cap A_1 = \emptyset$. Let $E_2 = E_1 \cup E(f_2)$ and let Γ_2 be the graph $(\mathcal{H} \cup V, E_2)$. Let C_2 be the connected component of Γ_2 containing H_0 . Choose a set A_2 in $[V]^k$ different from A_0 and A_1 , and let f_3 be a transversal of \mathcal{H} with $f_3[\mathcal{H}] \cap A_2 = \emptyset$.

Continuing in this way we obtain an ascending sequence of countable graphs $C_i, 1 \leq i < \omega$, sets $A_i \in [V]^k$, and transversals f_i of \mathcal{H} with $f_i[\mathcal{H}] \cap A_i = \emptyset$, so that whenever $f(H) \in V(C_i)$ there holds $H \in V(C_{i+1})$. Note that the choice of the sets A_i has been arbitrary so far. Writing

$V_0 = V \cap \bigcup_{i < \omega} V(C_i)$, we can choose the sets A_i in such a way that each member of $[V_0]^k$ is chosen as A_i for some $i < \omega$. Let $\mathcal{H}_0 = \bigcup_{i < \omega} V(C_i) \cap \mathcal{H}$. For each $H \in \mathcal{H}_0$ let $H' = H \cap V_0$. Let $\mathcal{H}'_0 = (H' : H \in \mathcal{H}_0)$. By our construction $\mathcal{H}'_0 - A$ has a transversal for every $A \in [V_0]^k$. Since \mathcal{H}'_0 is countable, it follows that it has a k -transversal, and hence \mathcal{H}_0 has a k -transversal contained in V_0 .

Let now $\mathcal{H}^0 = \mathcal{H} \setminus \mathcal{H}_0$, and let β_1 be the first ordinal such that $H_{\beta_1} \in \mathcal{H}^0$. Repeat the same construction with \mathcal{H}^0 and H_{β_1} , replacing \mathcal{H} and H_0 , respectively. We obtain a countable subfamily \mathcal{H}_1 of \mathcal{H}^0 , and a countable subset V'_1 of V , such that $H_{\beta_1} \in \mathcal{H}_1$ and \mathcal{H}_1 has a k -transversal into V'_1 . Define $V_1 = C_1 \setminus V_0$. Repeating this process until \mathcal{H} is exhausted yields the proof.

Proof of Corollary 1a. Suppose first that \mathcal{H} has a k -transversal. Then, by Theorem 1, $\mathcal{H} - S$ has a transversal for every $S \in [V]^k$. Therefore \mathcal{H} has a transversal. Let f be a string in V with $|\text{dom } f| \geq k$. Let $T = \{f(0), \dots, f(k-1)\}$ and f' be the string obtained from f by deleting its first k terms. Let q' be the function related to $\mathcal{H} - T$ as q is related to \mathcal{H} . Then $q'(f') \geq 0$ since $\mathcal{H} - T$ has a transversal, and so k applications of [6, Lemma 2.10] (whose proof remains valid for uncountable strings) yield $q(f) \geq k + q'(f') \geq k$.

Conversely, suppose that \mathcal{H} has a transversal and $q(f) \geq k$ for each string f with $|\text{dom } f| \geq k$. By Theorem 1 it suffices to show that $\mathcal{H}' = \mathcal{H} - A$ has a transversal for every $A \in [V]^k$. Since \mathcal{H}' has a transversal, it follows that in \mathcal{H}' for each string f there holds $q(f) \geq 0$. Let h be a transversal of \mathcal{H} . Then $\mathcal{H}' \setminus h^{-1}[A]$ has a transversal. By [1, Lemma 14] if a family \mathcal{G} has a countable subfamily \mathcal{F} such that $\mathcal{G} \setminus \mathcal{F}$ has a transversal and if in \mathcal{G} there holds $q(f) \geq 0$ for each string f then \mathcal{G} has a transversal. Thus \mathcal{H}' has a transversal, completing the proof.

Proof of Corollary 1c. Suppose that \mathcal{H} has a k -transversal. Then by Theorem 1 it has a transversal. Let \mathcal{C} be a nonempty subfamily of \mathcal{H} . Since \mathcal{H} has a k -transversal, it follows that \mathcal{C} has a k -transversal and so, by Theorem 1, we can select a set $S \in [\bigcup \mathcal{C}]^k$ and a transversal f of \mathcal{C} such that $S \cap f[\mathcal{C}] = \emptyset$, which shows that the inequality $|\bigcup \mathcal{C} \setminus f[\mathcal{C}]| < k$ is false for some transversal f of \mathcal{C} .

Assume that \mathcal{H} possesses the property in the statement and let f be a string with $|\text{dom } f| \geq k$. If $\mathcal{C} = \{H \in \mathcal{H} : H \subseteq \text{rng}(f)\}$ then by [6, Lemma 1.1] $|\bigcup \mathcal{C} \setminus h[\mathcal{C}]| \leq q(f)$ holds for every transversal h of \mathcal{C} . Now we can use Corollary 1a.

Proof of Theorem 3. By Theorem 1, if \mathcal{H} has a k -transversal then it has a transversal. If all members of \mathcal{H} are infinite, then $q(f) \geq \|\text{dom } f\|$ for each string f , and hence, by Corollary 1a, if \mathcal{H} has a transversal then it has a k -transversal.

ACKNOWLEDGMENT

The authors are grateful to the referee, whose suggestions improved the exposition at several points.

REFERENCES

1. R. AHARONI, On the equivalence of two conditions for the existence of transversals, *J. Combin. Theory Ser. A* **34** (1983), 202–214.
2. M. HALL, Distinct representatives of subsets, *Bull. Amer. Math. Soc.* **54** (1948), 922–928.
3. P. HALL, On representatives of subsets, *J. London Math. Soc.* **10** (1935), 26–30.
4. G. HOFFMAN AND P. KOMJÁTH, The transversal property implies property B, *Period. Math. Hungar.* **7** (1976), 179–181.
5. L. LOVÁSZ, A generalisation of König's theorem, *Acta Math. Acad. Sci. Hungar.* **21** (1970), 443–446.
6. C. ST. J. A. NASH-WILLIAMS, Another criterion for marriage in denumerable societies, *Ann. Discrete Math.* **3** (1978), 165–179.
7. K.-P. PODEWSKI AND K. STEFFENS, Injective choice functions for countable families, *J. Combin. Theory Ser. B* **21** (1976), 40–46.